

Isolation of lactic acid bacteria (LAB) species as probiotic from intestinal contents of common carp *Cyprinus carpio* L.
عزل بكتيريا العصيات اللبنية (معززات حيوية) من مكونات امعاء اسماك الكارب الشائع *Cyprinus carpio* L.

Jamal K.H.Al-Faragi

Sundus A.A.Alsaphar

Veterinary College/ Baghdad University

سندس عون علي

جمال خلف عطية الفراجي

كلية الطب البيطري / جامعة بغداد

Abstract

The presence of lactobacilli were investigated in the intestine of 50 fish specimens common carp *cyprinus carpio* weight rang between 400-1500g in weight from commercial farms in north of Baghdad through period of September to December 2010. All isolates were Gram positive ,catalase negative short rod usually non motile and non sporulating bacteria that produce lactic acid as a major or sole product of their fermentative metabolism.The obtained data showed that various species of lactic acid bacteria were found with high levels in total number about 10^4 - 10^7 CFU/g of intestinal content, physiological and biochemical characteristics of 8 strains isolated from intestine J1, J3, J4, J6, J7, J8, J9, J10 that can be categorized in two metabolic group facultative and obligate heterofermentation ,also all isolates were resistente to all antibiotic sensitivity OA2A -p Disc except isolates J1, J3, J4, J8, J9 were sensitive to erythromycin in concentration 60mcg .The aim of this study was to characterize of lactic acid bacteria isolated from the gastrointestinal tract of common carp *Cyprinus carpio* to be used as probiotic.

المستخلص

فحصت امعاء 50 سمكة نوع الكارب الشائع *Cyprinus carpio* بوزن يتراوح بين 400-1500 g المرربة في احواض شمال بغداد للمدة من ايلول الى كانون الاول عام 2010 عن وجود بكتيريا العصيات اللبنية اذ وجد ان جميع العزلات كانت بكتيريا موجبة لصبغة كرام عسوية قصيرة ,سالبة لفحص الكاتليز غير متحركة ولا تكون سبوروات ومنتجة حامض اللبنيك من عملياتها الايضية وبمعدل يتراوح بين 10^4 - 10^7 cfu/g من مكونات الامعاء وان الموصفات الفيزيائية والكيموحيوية بينت وجود 8 عزلات سميت J1 , J3 , J4 , J6 , J7 , J8 , J9 , J10 غير متجانسة اجبارية واختيارية التخمر , كانت جميع العزلات مقاومة للمضادات الحياتية في OA2A-P Disc ماعدا عزلة رقم J1 , J3 , J4 , J8 , J9 كانت حساسة للارثرومايسين بتركيز 60mcg هدفت الدراسة التحري عن العصيات اللبنية في القناة الهضمية في اسماك الكارب الشائع ودراسة مواصفاتها الفيزيائية والكيمو حيوية لغرض استعمالها كمعززات حيوية Probiotic لتحسين النمو والحالة الصحية ونسبة البقاء في الاسماك .

Introduction

Lactic acid bacteria are gram positive, non spore forming and catalase negative rods or cocci produce circular white and waxy colonies that ferment various carbohydrates mainly to lactate and acetate and various amino-acids can utilize but not usually reduce nitrate, Vitamins and minerals are essential for their growth [1]are commonly associated with nutritious environments like food. decaying material and the mucosal surfaces of the gastrointestinal and urogenital tract [1,2,3].Lactic acid bacteria are also part of normal intestinal flora of fish [4]. most of the evidences come

from salmonid species [4,5,6] Few studies have described lactic acid bacteria in other fishes [7,12]. [7] described the presence of lactic acid bacteria including lactobacillus in the intestines of various fish species at larvae fry and fingerling stages inhabiting ponds in Ukraine. They give information on the changes in their composition as a function of the season of the year and life stage of the fish .It was discussed that some human activities like artificial feeding of fish in ponds would have had an effect on the bacterial companion and load like common carp *Cyprinus carpio*» which showed the highest content of lactic acid bacteria in intestine

The distribution of lactic acid bacillus in the intestinal content of river fish and reported that various species of lacto bacilli were present in relatively high numbers in the intestines of edible fresh water fish from the river, specially in warm season but in low numbers in cold season[8] . [9]was pointed out studied the composition of intestinal lactic acid bacteria in carp fish and show that predominant LAB was *Lactococcus lactis* in July and *Lactococcus raffinolactis* in December and yearlong analysis of changes in LAB composition was performed in common carp from April 2002 to March 2003 . The predominant LAB was *L. lactis* in summer when water temperature was above 20 c° and *L. raffinolactis* in winter, when water temperature was ranges between 4-10°C.

The aim of present study was to characterization of LAB isolated from intestines of of the common carp « *Cyprinius carpio*» of commercial farms near Baghdad.

Materials and methods

Fish bacterial media and reagents

Fish

Several groups of adults common carp were obtained from commercial fish farms in the north of Baghdad from September 2010- to December 2010 and maintained in aquarium filled with tap water at 21 c° .

Chemicals and media

Analytical grade chemicals and dyes were obtained from Al-Kindi company for production of veterinary vaccines and drugs , bacteriological media were obtained from oxoide UK which include set of biochemical media, blood agar base,,Nutreint agar , gas generating Kit and sens test disk from lamb GT. Manchester; England finally MRS media were obtained from Himedia ,India .

Isolation of bacterial strain

Various samples of the common carp « *Cyprinus carpio*» of commercial farm in the north of Baghdad (average between 400-1500 g in weight). They were brought to the laboratory alive and sacrificed. The abdomen surfaces were thoroughly scrubbed with an alcohol (70% ethanol) and aseptically dissected to remove the intestines. The intestines were opened by longitudinal incision and thoroughly flushed with sterilized chilled normal saline solution (NSS) to remove feed materials, dirt and other impurities. The intestines were weighted macerated with sterile glass rod and homogenized in sterile NSS (1:10 wet/vol) using a vortex mixer. Samples of the thoroughly macerated and homogenized intestines were serially diluted in NSS and aseptically plated by pour plate technique on MRS agar and incubated anaerobically at 30° C for 2-3 days.

Well isolated colonies were picked up less than 10 colonies per each plate and all the samples were counted according to the method described by [10] and transferred to MRS broth. They were propagated twice and streaked on MRS agar to check the purity of the isolates and then stored in MRS soft agar 0.5% overlaid with 50% glycerol at -20 °C and pure culture were freeze dried and stored at -20 °C the isolates maintained in frozen stocks were propagated twice in MRS broth and used for further study, these were inoculated in to MRS broth and incubated at 30 °C for 18 h.

Characterization and Differentiation of LAB

Isolated strains were gram stained and examined microscopically for cellular morphology and Gram-stain phenotype, catalase activity was tested by spotting colonies with 3% hydrogen peroxide. Growth was assayed in MRS broth at 4, 10, 25, 30, 35 and 45 °C as well as pH 4.0, 5.0, 6.5, 8.5 and 9.0 incubated at 30 °C. Salt tolerance was tested with 6.5%, 10, 15% w/v NaCl in MRS broth production of acid and CO₂ from glucose was tested in MRS broth containing Darhams tube with citrate omitted [1] production of Ammonia in MRS broth omitting glucose and meat extract while containing 0.3% Arginine and 0.2% Sodium citrate replacing ammonium citrate was monitored using Nessler's reagent. Assays for gelatin hydrolysis and nitrate reduction [11]. Ability to ferment various carbohydrates was evaluated using MRS broth supplemented with filter sterilized sugar solutions to final concentration of 1% w/v and 0.004% phenol red without glucose and meat extract.

Antibiogram of LAB isolates

The isolates were inoculated into MRS broth individually and incubated for 24 hr about 25 ml of MRS agar was seeded with the cultures of LAB isolates 10⁶ CFU/ml mixed well. Poured in to sterile petriplates and stored at 4 °C for 1hr to solidify the media (OA21-P) antibiotics in a single ring were placed upside down pressed on the top of the agar plates and kept again at 4 °C for 1hr the plate were incubated at 30 °C for 24 hr. and 48hr resistance was defined as the absence of a growth inhibition zone around the discs.

Results and Discussion

Intestinal content of all fish sample analyzed for the presence of lactobacilli. Counts of intestinal lactic acid bacteria for common carp detected at the range of 10⁴–10⁷ CFU/g found in table (1) these results were supported by [12][13]; they estimated lactic acid bacteria counts 10⁷ to 10⁸ CFU/g in the intestine of common carp, while [14] showed the count of intestinal lactobacillus spp for Persian sturgeon and beluga were detected at the range of approximately 10^{5.3} to 10^{6.4} CFU/g respectively but [6] found the count of LAB on farm fish was 10⁰ -10³ CFU/g the reasons of all these different result were explained the effect of fish species, life stage of the fish, some human activities like artificial feeding in ponds and also season of the year [9] All the 8 isolates bacteria were found Gram positive while the morphology was coccobacilli arranged in small chain contain 2-6 cells or as a single tetrad and paired cell physiological behavior of all isolates is present in Table(2).

Table (1): Count of LAB isolates according to date and weight of sample common carp *Cyprinus carpio*

Sample number	Isolates	Date	Avareg weight of sample	Morphology of LAB isolation	Count of LAB isolation per gram
1	J1	September	750-850	Coccobacilli	9x10 ⁶ cfu /g
2	nil	September	500-650	nil	nil
3	J3	October	400-500	Coccobacilli	7x10 ⁷ cfu/g
4	J4	October	400-550	Coccobacilli	3.5x10 ⁷ cfu/g
5	nil	October	450- 600	nil	nil
6	J6	October	500- 740	Coccobacilli	2X10 ⁷ cfu/g
7	J7	October	600- 750	Coccobacilli	9x10 ⁶ cfu/g
8	J8	November	1000-1350	Coccobacilli	3x10 ⁵ cfu/g
9	J9	December	1200-1500	Coccobacilli	7x10 ⁴ cfu/g
10	J10	December	1100-1150	Coccobacilli	1x10 ⁴ cfu/g

Table (2): Physiological properties of lactic acid bacteria isolates.

Test	Isolates of Lactic acid bacteria							
	J1	J3	J4	J6	J7	J8	J9	J10
Morphology	Strept Cocco bacilli	short rod	Tetrad short rod	short rod	Single paired short rod	Tetrad coccobacilli	short rod	Tetrad coccobacilli
Growth at 4	-	-	-	-	-	-	-	-
Growth at 10	+	+	+	+	+	+	+	+
temperatur e 25	++	++	++	++	++	++	++	++
Temperatu re 30	+++	+++	+++	+++	+++	+++	+++	+++
35	+++	++	++	+++	++	++	++	++
45	+++	-	-	-	-	++	++	-
	++							
Growth at pH 4	+	+	+	+	+	+	+	+
5	++	++	++	++	++	++	++	++
6.5	+++	+++	+++	+++	+++	+++	+++	+++
8.5	+	+	-	+	+	+	+	+
9	-	-	-	-	-	-	-	-
Growth at Nacl% 5	+	+	+	+	+	+	+	+
6.5	+	-	-	-	+	+	+	-
10	W	-	-	-	-	W	-	-
15	-	-	-	-	-	-	-	-

(+)growth (-) no growth (w) weak growth

Luxurious growth was observed for all 8 isolates at 30 and 35C° while weak growth was observed at 10 C° and no growth was observed at 45 C° for isolates J3, J4, J6, J7, L10.

The results are shown in the same table; effect of different pH , Nacl concentration (salt tolerance) on the growth Biochemical as well as carbohydrate fermentation and behavior of these isolates against various antibiotics were also found in table (3,4,5) .

Table (3): Biochemical properties of LAB isolate

Test	Bacteria isolates							
	J1	J3	J4	J6	J7	J8	J9	J10
Gram stain	G+ev	G+ev	G+ev	G+ev	G+ev	G+ev	G+ev	G+ev
Motility	-	-	-	-	-	-	-	-
Spore forming	-	-	-	-	-	-	-	-
Catalase	-	-	-	-	-	-	-	-
Oxidase	-	-	-	-	-	-	-	-
Gelatin hydrolysis	-	-	-	-	-	-	-	-
Starch hydrolysis	-	-	-	-	-	-	-	-
CaCO ₃ 0.8% lyses	+	+	+	+	+	+	+	+
CO ₂ production from glucose	-	-	-	-	-	+	+	+
Nitrite reduction	-	-	-	-	-	-	-	-

(+) growth (-) no growth

Table(4): Carbohydrate Utilization of LAB isolates

Carbohydrates	Bacteria isolates							
	J1	J3	J4	J6	J7	J8	J9	J10
Glucose	+	+	+	+	+	+	+	+
Trehalose	+	+	+	+	+	+	+	+
Dextrine	+	+	+	-	+	+	+	+
Inuline	+	+	+	+	+	-	+	+
Lactose	+	+	+	+	+	+	+	+
Sorbitol	+	+	+	+	+	+	+	+
Adintol	+	+	+	+	+	+	+	+
L-arabinos	-	-	-	N.R	-	-	-	-
Sucrose	+	+	+	+	+	-	+	+
Dolcitol	+	+	+	+	+	+	+	+
Galactose	+	+	+	+	-	-	+	-
Sorbitos	-	-	-	-	+	-	+	+
Raffinos	+	+	+	+	+	+	+	+
Cellulose	-	-	-	-	+	-	-	+
Mannitof	+	+	+	+	+	+	-	+
Ffructose	+	+	+	+	+	+	+	+
Mannose	+	+	+	+	+	+	+	+
Maltose	+	+	+	+	+	+	+	+

(+) ferment (-) Not ferment (+*) delayed ferment more than 24 h
(NR) no reaction

On the basis of morphological, physiological and biochemical characters as well as sugar utilization all isolates identified were to the genus *Lactobacillus* but could not be assigned to any particular species by these characters it is interesting to note that the majority of lactic acid bacteria spp that have been isolated from fish were those species which were commonly found on meat animals and human [1]. [14] reported that lactobacilli found in their study were relatively similar to the species described [8] These authors reported *L. alimentarius*, *L. coryneformis*, *L. casei*, *L. sakei*, *L. pentosus*, *L. plantarum*, *L. bevis* and *L. oris*, as lactobacilli presented in the intestinal content of studied fish while [15] found the occurrence of typical lactobacilli as described by [1] were rare in fish this signified the need for proper classification for lactic acid bacteria from fresh water fish

In this study Most *Lactobacillus* examined had the capacity to ferment lactose and galactose. Most lactobacilli are able to ferment lactose by uptake of this disaccharide by a specific permease and splitting it by *s. galactosidase* for further phosphorylation of galactose and glucose [1]. Because lactose is only present in milk and milk derivatives, it is possible that these strains have evolved from environments related with mammals, as was suggested for other lactose positive *Lactobacillus* [16]. Lactose may be present or was present in the environment as a waste resulting from livestock production and disposal effluents from dairy factories another component often fermented by the strains was the amino-sugar N- acetyl – glucosamine. a compound present in peptidoglycans, in blood, chitin and as one of the main constituents of mucus in the gastrointestinal tract [17] The carbohydrate and protein constitutes above 40% of the weight of the mucus [18] or higher values [17].

Fish at all life stages may expose to the bacteria from the environment some of them are detrimental and others are beneficial. Current methods for control of pathogens in the fish farms should be improved by studying the beneficial bacteria. As *Lactobacillus* has many documented health effect [19]. And naturally present in the gastrointestinal tract of man and animals [20] we started studies aimed to investigate the intestinal *Lactobacillus* in fish with the goal of selecting a strain to be used as a feed supplement for warm fresh water fish. Knowledge on the presence of *Lactobacillus* as a natural flora in fish may lead to further application to improve to improve growth performance, survival rate and healthy of fish.

References

1. Kandler, O and Weiss, N. (1986). Genus *Lactobacillus* Beijerinck (901, 212). In: Sneath, P. H. A.; Mair, N. S.; Sharpe, M. E. and Holt, J. G. (eds.) Bergey's manual of systematic bacteriology. V. 2, Baltimore: Williams and Wilkins, 1209-1234.
2. Havenaar, R.; Ten Brink, B. and Huis in Veld, J. H. J. (1992). Selection of strains for probiotic use. in: Fuller, R. (ed), probiotics: The scientific basis (1st Edn.), London Chapman and Hall, 209-224.
3. Walstra, P.; Geurts, T. J.; Noomen, A.; Jellema, A. and van Boekel, M. A. J. S. (1999). Dairy technology, principles of milk properties and processes. New York, Marcel Dekker, Inc., 727.
4. Ringo, E. and Gatesoupe, F. J. (1998). Lactic acid bacteria in fish: a review. *Aquaculture*, 160:177-203.

5. Ringo, E. and Olsen, RE. (1999) .The effect of diet on aerobic bacterial Flora associated with intestine of Arctic charr (*Salvelinus alpinus* L.). J.Appl. Microbiol.,86 :22-28.
6. Ringo, E.; Bendiksen, H.; Wesmajervi, M.; Olsen, R.; Jansen, R. and Mikkelsen , H. (2000) . Lactic acid bacteria associated with the digestive tract of atlantic Salmon (*Salmo salar* L.) J. Appl . Microbiol., 89: 317- 322.
7. Kvasnikov, El;Kovalenk,NK and Materinskaya, L.G .(1977) . Lactic acid bacteria of water fresh fish microbiology, 46:619-624.
8. Bucio Galindo, A; Hartemink , R; `Schrama , JW; Verreth , JAJ. and Rombouts , FM.(2006).Presence of lactobacilli in the intestinal content of fresh water fish from a river and and from a farm with recirculation system .Food Microbiol., 23:476-482.
9. Tatsuro Hagi, D. T. ; Yasutada L. and Takayuki, H. (2004) .Diversity and seasonal changes in lactic acid bacteria in the intestinal tract of cultured fresh water fish . Applied Biochemistry, University of Tsukuba, 1-1-1 Ten – Nodai Tsukuba, Ibaraki: 305-8572.
10. Thapa, N; Pal, J .and Tamang, JP.(2006). Phenotypic identification and Technological Properties of lactic acid bacteria isolated from traditionally processed fish products of the Eastern Himalayas. Int.J. food Microbiol., 107 ;33-38.
11. Hammes,WP.and Vogel, RF. (1995). The genus lactiobacillus. In: wood, BJB and Holzapfel, WH (eds), The lactic acid bacteria. Vol. 2, London, Blackie Academic. and Professional: 19-54 .
12. Cai. Y; Suyanandana .P; Saman, P.and Benno,Y. (1999). Classification and characterization of lactic acid bacteria isolated from the intestines of common carp and water fresh prawns. J .Gen .Appl . Microbiol ., 45 (4):177-184.
13. Khali, A; TKmechi, A.and Agh,N. (2009). Characterization of lactic Acid bacteria isolated from the intestines of common carp of west Azarbaijn, Iran. Joural of Animal and Veterinary Advances. Vol 8 (6): 1162- 1164.
14. Ghanbari , M. ; Rezaci , M. ; Jami , M.; and Nazari , R. M .(2009).Isolation and characterization of lactobacillus species from intestinal contents of beluga (*Huso huso*) and Persian sturgeon (*Acipenser Persicns*).Iranian Journal of Veterinary Research , Shiraz Uuiversity , 10(2)Ser . No. 27.
15. Parvath, S. N. and Puthuvallil, K. S.(2005). Biochemical Characterization of Lactic acid Bacteria isolated from fish and prawn. Journal of Culture Collections. 48-52.
16. Garvie, EI.(1984). Taxonomy and Identification of diary bacteria.in: Davies, FL .and Law , BA .(eds) Advanas in the microbiology and biochemistry of cheese and fermented milk. London: Elsevier Applied Science Pubishers, 35-65.
17. Hicks, SI.; Theodoropoulos , G.;Garrington , SD and Corfield , AP. (2000). The role of mucins in host-parasite interactions. Part 1- Protozoam Parasites. Parasitol Today .16 : 476 – 481.
18. Stephen, AM. (1985). Effect of food on the intestinal microflora. In: Hunter, JO.and Alun, J. V.(eds.), Food and the gut. Sussex, England, Bailliere Tindall., 57-77.
19. Onwehand, AC.;kirjavainen , PV. ; Gronlund, MM.; Isolauri, E.and Salminen, S.(1999).Adhesion of probiotic micro-organisms to intestinal mucus. Int. Dairy. J,9:623-630.
20. Fuller, R.(1989). Probiotics in man and animals. J. Appl. Bacteriol., 66:365 – 378.